

Realizing the chromatic numbers and orders of spinal quadrangulations of surfaces

Serge Lawrencenko

(Department of Higher Mathematics 1, National Research
University of Electronic Technology, Zelenograd, Russia
E-mail: lawrencenko@hotmail.com)

July 8, 2012

Abstract

A method is suggested for construction of quadrangulations of the closed orientable surface with given genus g and either (1) with given chromatic number or (2) with given order allowed by the genus g . In particular, N. Hartsfield and G. Ringel's results [Minimal quadrangulations of orientable surfaces, J. Combin. Theory, Series B 46 (1989) 84–95] are generalized by way of generating minimal quadrangulations of infinitely many other genera.

Keywords: embedding of graph, coloring of graph, Betti number, minimal quadrangulation, computer animation.

MSC Classification: 05C10 (Primary); 05C15, 05C75, 57M15 (Secondary).

1 Introduction

The purpose of this note is to suggest a method for generating quadrangulations of closed orientable surfaces with given properties. Those surfaces arise as the thickenings of spatial graphs called the spines. The operation of thickening a given graph G builds a surface around G which is used in computer animation. See [1]. The first step of thickening G consists in building a cylinder around each edge of G by setting four quadrilaterals around that edge. Therefore a thickening of a spine graph is represented by its so-called spinal quadrangulation. The properties of the spinal quadrangulation are determined by its spine. In particular, one can control the

genus, chromatic number, and order of the quadrangulation by choosing a suitable spine. Hence, the question arises of how wide the spectra of these three parameters could be.

2 Spinal quadrangulations

If a (finite, undirected, simple) graph G is 2-cell embedded in the sphere with g handles, Σ_g , the components of $\Sigma_g - G$ are called *regions*. A *quadrangulation of Σ_g with graph G* is a 2-cell embedding $G \hookrightarrow \Sigma_g$ in which each region is bounded by a simple circuit of length 4 in G . The *chromatic number* $\chi(G)$ of G , as well as of any quadrangulation with this graph, is the minimum number of colors sufficient for coloring the vertices of G so that adjacent vertices receive different colors. The (*first*) *Betti number* of a connected graph G is given by $\beta(G) = |E(G)| - |V(G)| + 1$, where $|V(G)|$ and $|E(G)|$ stand for the cardinalities of the vertex and edge sets of G (respectively), and $|V(G)|$ is also called the *order* of G . Especially, for the complete graph K_n of order n , $\beta(K_n) = \frac{1}{2}(n-1)(n-2)$. The *2-fold interlacement* of G is denoted by $G[:]$ and is defined to be the graph which has vertex set $V(G[:]) = V(G') \cup V(G'')$, where G' and G'' are two disjoint copies of G , and as edges the set $E(G[:]) = E(G') \cup E(G'')$ plus the edges that join each vertex $v' \in V(G')$ to each vertex in G'' adjacent to the corresponding vertex $v'' \in V(G'')$ (but not to v'' itself).

Lemma 2.1. (White [3], Craft [4]) *For any non-trivial connected graph G , there exists a quadrangulation $G[:] \hookrightarrow \Sigma_{\beta(G)}$.*

Any quadrangulation $G[:] \hookrightarrow \Sigma_{\beta(G)}$ is called a *spinal quadrangulation* with *spine* G and *genus* $g = \beta(G)$. By Lemma 2.1, the genus of a spinal quadrangulation is equal to the Betti number of the spine. The following theorem is a quadrilateral analogue of the result of [2] on triangulations.

Theorem 2.2. *For any integers $g \geq 0$ and $n \geq 2$ such that $g \geq \beta(K_n) = \frac{1}{2}(n-1)(n-2)$, there exists a spinal quadrangulation of Σ_g with chromatic number n .*

Proof. Since it is obvious that $\chi(G[:]) = \chi(G)$, it follows from Lemma 2.1 that we can use as a spine any graph G with $\beta(G) = g$ and $\chi(G) = n$. To construct such a spine, we start up with K_n . If $\beta(K_n) = g$, we're done. If $\beta(K_n) < g$, we take the ladder graph with $g - \beta(K_n)$ linearly independent cycles (of length 4) and glue one of its ends on K_n . $\square$

The spectrum $\{n\}$ described in Theorem 2.2 for possible chromatic numbers is full for any fixed genus $g \geq 0$. In fact, $\frac{1}{2}(\chi(G) - 1)(\chi(G) - 2) = \frac{1}{8}(2\chi(G) - 3)^2 - \frac{1}{8} \leq \beta(G)$, where the inequality is provided by [5, 6], whence the maximum chromatic number n of a graph with given Betti number $g \geq 0$ is attained by the largest complete graph K_n such that $\beta(K_n) \leq g$.

Yet, the question remains of what the limitations of the proposed approach are when speaking about the whole spectrum of possible chromatic numbers for a quadrangulation of a given surface. If K_n has a quadrilateral embedding in Σ_g (for instance when $n \equiv 5 \pmod{8}$), then $g = 1 + \frac{1}{8}n(n - 5)$ and one may expect that for each h such that $2 \leq h \leq n$ there would be a quadrangulation of Σ_g with chromatic number h . Meanwhile, the largest integer m such that

$$\beta(K_m) = \frac{1}{2}(m - 1)(m - 2) \leq g = 1 + \frac{1}{8}n(n - 5)$$

is $m = \lfloor \frac{1}{2}(3 + \sqrt{8g + 1}) \rfloor \approx \frac{n}{2}$ for large n . Therefore, by using quadrilateral embeddings of spine graphs, we manage to cover about a half of possible values of the chromatic number.

3 Minimal quadrangulations

We start up with two corollaries of Lemma 2.1.

Corollary 3.1. *For any integer $p \geq 2$ there exists a quadrangulation $K_p[:]$ $\hookrightarrow \Sigma_g$ with $g = \beta(K_p) = \frac{1}{2}(p - 1)(p - 2)$.*

Corollary 3.2. *Let p be an integer ≥ 3 and let $(K_p - e)$ be K_p minus one edge. Then there exists a quadrangulation $(K_p - e)[:]$ $\hookrightarrow \Sigma_g$ with $g = \beta(K_p) - 1$.*

A quadrangulation of a fixed surface Σ_g is said to be *minimal* in Σ_g if the number of vertices is minimal among all (not necessarily spinal) quadrangulations of Σ_g . The quadrangulations of Corollaries 3.1 and 3.2 were first discovered (in terms of general octahedral graphs) by Hartsfield and Ringel [7], who also showed their minimality in Σ_g whenever $p \geq 4$ for Corollary 3.1, and $p \geq 8$ for Corollary 3.2. These corollaries are special cases of our main theorem, stated shortly, and correspond to the particular cases $m = 0$ and $m = 1$, respectively. Using the method of current graphs for $m \geq 2$ would have been very complicated, so Hartsfield and Ringel had to stop at $m = 1$. In contrast, the spinal method enables generalization of the results of [7] to an arbitrary m . This demonstrates one way in which the spinal method is useful.

Theorem 3.3. *Let $(K_p - me)$ be K_p minus m edges ($0 \leq m \leq \beta(K_p)$). If $(K_p - me)$ is connected, there exists a quadrangulation $(K_p - me)[:] \hookrightarrow \Sigma_g$ with $g = \beta(K_p) - m$. Moreover, any such quadrangulation is minimal in Σ_g with $g \geq 1$ whenever $p \geq 4(m + 1)$.*

Proof. Compute $g = \beta(K_p - me) = \frac{1}{2}(p - 1)(p - 2) - m$, and the existence follows from Lemma 2.1. It remains to prove minimality of the constructed quadrangulation. Let $\alpha_0, \alpha_1, \alpha_2$ denote the number of vertices, edges, and regions of an arbitrary quadrangulation of Σ_g . By Euler's equation, we have $\alpha_0 - \alpha_1 + \alpha_2 = 2 - 2g$. Furthermore, since any pair of vertices are joined by at most one edge, we have $\alpha_1 \leq \binom{\alpha_0}{2}$, and since each edge meets exactly two regions, we have $4\alpha_2 = 2\alpha_1$. From these it can be derived that $\alpha_0^2 - 5\alpha_1 + (8 - 8g) \geq 0$. This quadratic inequality has the solution:

$$\alpha_0 \geq \left\lceil \frac{1}{2} \left(5 + \sqrt{32g - 7} \right) \right\rceil. \quad (1)$$

With g computed in the beginning of the proof, we find

$$32g - 7 = 16p^2 - 48p + 25 - 32m. \quad (2)$$

Now, since the constructed quadrangulation has $2p$ vertices, it follows from Eqs. (1) and (2) that it is minimal in Σ_g whenever the following double inequality holds:

$$2p - 1 < \frac{1}{2} \left(5 + \sqrt{16p^2 - 48p + 25 - 32m} \right) \leq 2p.$$

This can be rewritten as $16p^2 - 56p + 49 < 16p^2 - 48p + 25 - 32m \leq 16p^2 - 40p + 25$, or as

$$\begin{cases} 8p > 24 + 32m, \\ 8p \geq -32m. \end{cases}$$

Since the second inequality is a tautology, we conclude that $p > 3 + 4m$, hence $p \geq 4 + 4m$. The proof is complete. $\square$

Theorem 3.3 provides many new minimal quadrangulations for infinitely many genera g not covered by Hartsfield and Ringel [7]. For example, the quadrangulations $K_{12}[:] \hookrightarrow \Sigma_{55}$, $(K_{12} - e)[:] \hookrightarrow \Sigma_{54}$, and $(K_{12} - 2e)[:] \hookrightarrow \Sigma_{53}$ are minimal in the corresponding surfaces (and have 132, 130, and 128 regions, respectively). The first two are covered by [7] (or Corollaries 3.1 and 3.2), but the minimal quadrangulation of Σ_{53} is a new one. Note that a quadrangulation is minimal in the sense of [7] if it has the minimum number of regions, but since the definition assumes the surface is fixed, their definition agrees with the one given in the beginning of this section.

The following is an analogue of Theorem 2.2.

Corollary 3.4. *For any integers $g \geq 0$ and $p \geq 2$ such that $g \leq \beta(K_p)$, there exists a spinal quadrangulation of Σ_g with order $2p$.*

Proof. The order of a spinal quadrangulation is twice the order of its spine. By Theorem 3.3, we can take $(K_p - me)$ as a spine, letting m be $(\beta(K_p) - g)$. Clearly, it is possible to remove this number of edges from K_p so that the remaining graph is still connected. $\square$

The spectrum $\{2p\}$ described in Corollary 3.4 for possible orders is full for fixed genus; there are infinitely many spinal quadrangulations of arbitrarily large even order in each surface Σ_g . Solving the quadratic equation $\beta(K_p) = g$ for p , we come to the following formula for the order of a minimal spinal quadrangulation with genus $g \geq 0$:

$$V_{\min}^{(s)}(g) = 2 \left\lceil \frac{1}{2}(3 + \sqrt{8g + 1}) \right\rceil. \quad (3)$$

We'll shortly provide a partial formula for $V_{\min}(g)$ —the order of a minimal general quadrangulation with genus g . Clearly, $V_{\min}^{(s)}(g) \geq V_{\min}(g)$. For example, $V_{\min}^{(s)}(0) = V_{\min}(0) = 4$ by Eq. (3). It should be noted that Hartsfield and Ringel [7] assert that $V_{\min}(0) = 8$ because their definition of a quadrangulation, in comparison to the definition given in the beginning of Section 2, has an additional requirement as follows: the intersection of any two distinct regions is either empty or at most one edge and at most three vertices. However, any minimal quadrangulation in the sense of our definition has no vertices of degree 2 whenever $g \geq 1$, and therefore satisfies this requirement. By Eq. (3), $V_{\min}^{(s)}(1) = 6$ and $V_{\min}^{(s)}(2) = 8$, whereas it is shown in [7] that $V_{\min}(1) = 5$ and $V_{\min}(2) = 7$.

Corollary 3.5. *For $g \geq 3$, $V_{\min}(g) = 2\lceil a(g) \rceil$ whenever $\lceil a(g) \rceil = \lfloor b(g) \rfloor$, where $a(g) = \frac{1}{2}(3 + \sqrt{8g + 1})$ and $b(g) = \frac{1}{4}(7 + \sqrt{32g - 15})$.*

Proof. It follows from Theorem 3.3 that $V_{\min}(g) = V_{\min}^{(s)}(g) = 2p$ whenever this double inequality holds: $\beta(K_p) + 1 - \frac{p}{4} \leq g \leq \beta(K_p)$ or, equivalently, $a(g) \leq p \leq b(g)$. Note that whenever $g \geq 3$ (and so $p \geq 4$) we have $a(g) \leq b(g)$, and also observe that the closed interval $[a(g), b(g)]$ has length < 1 and can contain at most one integer, $\ell(g)$. Furthermore, such an $\ell(g)$ exists if $\lceil a(g) \rceil = \lfloor b(g) \rfloor (= \ell(g))$, in which case we have $V_{\min}(g) = V_{\min}^{(s)}(g) = 2\ell(g)$. $\square$

The formula of Corollary 3.5 applies to infinitely many values of g . For example, for each $p \geq 4$ by letting $g = \beta(K_p)$ we find $a(g) = p (= \ell(g))$, and so $V_{\min}(g) = 2p$.

Acknowledgments: I'm grateful to Vladimir P. Korzhik for his constructive feedback and comments. I'm also grateful to Thom Sulanke who endorsed me for arXiv.org.

References

- [1] S. Brandel, D. Bechmann, Y. Bertrand, The thickening: an operation for animation, *J. Visualization & Computer Animation* 11 (2000) 261–277.
- [2] F. Harary, S. Lawrencenko, V. Korzhik, Realizing the chromatic numbers of triangulations of surfaces, *Discrete Math.* 122 (1993) 197–204.
- [3] A. T. White, On the genus of the composition of two graphs, *Pacific J. Math.* 41 (1972) 275–279.
- [4] D. L. Craft, On the genus of joins and compositions of graphs, *Discrete Math.* 178 (1998) 25–50.
- [5] W. C. Coffman, S. L. Hakimi, E. Schmeichel, Bounds for the chromatic number of graphs with partial information, *Discrete Math.* 263 (2003) 47–59.
- [6] G. Chartrand, P. Zhang, *Chromatic Graph Theory*, Chapman & Hall/CRC Press, Boca Raton, FL, 2009.
- [7] N. Hartsfield, G. Ringel, Minimal quadrangulations of orientable surfaces, *J. Combin. Theory, Series B* 46 (1989) 84–95.